

IMPACTS OF RECENT INVASIVE SPECIES ON NEARSHORE FISHES

**John M. Dettmers², Charles P. Madenjian, Paul J. Allen, Steven A.
Pothoven, and Thomas F. Nalepa**

Introduction

Invasive species have impacted Lake Michigan's fisheries for 70 years. The passage through the Welland Canal by sea lamprey, after it invaded Lake Ontario, led to its establishment in Lake Michigan where it was first seen in 1936 (Christie and Goddard 2003). A combination of overfishing and sea lamprey predation led to the extirpation of lake trout (Coble et al. 1990; Hansen 1999) and to the extirpation or extinction of several deepwater ciscoes endemic to the Great Lakes (Coon 1999). Alewives, probably through interference with reproduction, likely caused the decline in abundances of deepwater sculpins and yellow perch during the 1960s and may have delayed the recovery of burbot in Lake Michigan until the 1980s (Madenjian et al. 2002). From 1959 to 1999, the rate of new introductions into the Great Lakes has increased to more than one species per year

J.M. Dettmers. Illinois Natural History Survey, Lake Michigan Biological Station, 400 17th Street, Zion, IL, 60099, U.S.A. (Present address: Great Lakes Fishery Commission, 2100 Commonwealth Blvd., Suite 100, Ann Arbor, MI, 48105, U.S.A.).

C.P. Madenjian. U.S. Geological Survey, Great Lakes Science Center, 1451 Green Road, Ann Arbor, MI, 48105, U.S.A.

P.J. Allen. Aquatic Biology and Fisheries Center, Department of Biology, Ball State University, Muncie, IN, 47306, U.S.A.

S.A. Pothoven. Cooperative Institute of Limnology and Ecosystem Research, University of Michigan/Great Lakes Environmental Research Laboratory, 1431 Beach Street, Muskegon, MI, 49441, U.S.A.

T.F. Nalepa. National Oceanic and Atmospheric Administration, Great Lakes Environmental Research Laboratory, 2205 Commonwealth Blvd., Ann Arbor, MI, 48105, U.S.A.

² Corresponding author (e-mail: jdettmers@glfc.org).

(Grigorovich et al. 2003), resulting in the establishment of numerous invaders during the past 20 years. In this report, we focus on (1) the status of six recent invasive species during 2000-2004, (2) the identity of other species that could invade in the near future, (3) examples of their impacts on the nearshore fish community, and (4) the implications for fishery management.

Status

Spiny Water Flea

Bythotrephes longimanus was first recorded in Lake Michigan in 1986 (Evans 1988). Native to the Ponto-Caspian region of Europe, this predatory cladoceran has been implicated in the reduction of small-bodied daphnids since the 1980s (Lehman 1991). Although few regular estimates of lakewide densities are available, mean density of *Bythotrephes* averaged across eight sites was 570/m² in offshore waters during September 2000 (Pothoven et al. 2003). Except for being implicated in reduced diversity of the offshore *Daphnia* species complex (Lehman and Cáceres 1993), this invader has had little apparent impact on nearshore fishes.

Zebra and Quagga Mussels

Zebra mussels were first documented in Lake Michigan in 1989 (Marsden 1992) and rapidly increased their population size in nearshore rocky habitats (Marsden 1992). Quagga mussels were first documented in Lake Ontario (May and Marsden 1992) and were present in Lake Michigan by 1997 (Nalepa et al. 2001). Although little quantitative monitoring of zebra mussels is conducted in Lake Michigan, an estimate of lakewide biomass is available from U.S. Geological Survey fall bottom-trawl assessments. Between 1999 and 2004, lakewide biomass of zebra and quagga mussels ranged from 14 kt (1,000s of metric tonnes) in 1999 to a peak of 43 kt in 2001 and returned to 14 kt in 2003 (Madenjian et al. 2005a). These lakewide estimates are likely conservative because trawls are not fished over the hard substrate on which zebra mussels prefer to attach (Fleischer et al. 2001; Coakley et al. 2002). Zebra mussel biomass was not separated from quagga mussel biomass in these trawl surveys, but observations suggest that the proportion of quagga mussels has been increasing in recent years.

Round Goby

This invader was first reported from Lake Michigan in 1993 (Charlebois et al. 1997). Since then it has spread slowly through southern Lake Michigan and most likely was transported via commercial shipping to ports farther north in the lake (Mills et al. 1993; Jude 2001), where it has further spread northward and eastward (Clapp et al. 2001). As round gobies spread into the lake, assessment trawls documented their increased relative abundance. Trawling by both the Michigan Department of Natural Resources (DNR) at Grand Haven and by Ball State University (BSU) in Indiana waters first detected round gobies in 1997 (Michigan DNR) and 1998 (BSU). Catch rates rose by two orders of magnitude within two years and, since then, have generally remained at similarly high levels (Lauer et al. 2004).

Fishhook Water Flea

Cercopagis pengoi, the fishhook water flea, was first recorded in Lake Michigan in 1999 (Charlebois et al. 2001). Like *Bythotrephes longimanus*, its relative, this invasive predatory cladoceran from the Ponto-Caspian region feeds on zooplankton. Unlike its relative, *Cercopagis* feeds primarily on smaller zooplankton, including rotifers, juvenile copepods, and small cladocerans (Vanderploeg et al. 2001; Benoit et al. 2002; Laxson et al. 2003). The abundance of this species, although variable, has trended upwards (Witt et al. 2005). Its impacts on the food web remain uncertain. Improved monitoring of its population dynamics and community impacts remains important to fully understand whether this invader will exert strong impacts on the nearshore food web.

Ruffe

Ruffe, a Eurasian fish, was first documented in Lake Michigan at Little Bay de Noc in 2002. It was found there again in 2003 and 2004 and since has spread to Big Bay de Noc. In other targeted and routine sampling around the lake, no other populations of ruffe have been reported. It is unclear why this species has expanded relatively slowly after introduction into Lake Michigan. The pattern of slow spread observed in Lake Superior, about 200 miles in eight years from 1987 to 1994 (Gunderson et al. 1998), may result from the apparent negative relationship between ruffe abundance and system productivity (Ogle 1998). Ruffe may initially colonize a location via commercial shipping but, once established, do not aggressively immigrate

into new areas. This behavior may provide a control method if ports are at risk.

Species on the Doorstep

Other potential invaders could arrive during the next few years. Of special concern is the possibility that silver carp or bighead carp, collectively known as Asian carp, could enter Lake Michigan through the Chicago Sanitary and Ship Canal (CSSC), the live food trade, or by other means. A permanent electric barrier is being constructed in the CSSC to deter movement across this artificial connection between the Mississippi River and Great Lakes drainages. Similarly, efforts among U.S. and Canadian agencies and legislative bodies are seeking to eliminate the trade in live Asian carp.

The northern snakehead is another potential invader. This species escaped into the Potomac River basin, most likely from aquarium releases. Specimens have been collected by the Wisconsin DNR and Michigan DNR on the non-Great Lakes waters of these states. One snakehead was collected by an angler while fishing in a Chicago harbor in October 2004. Based on an intensive sampling effort in the harbor, best estimates suggest that this snakehead was released from an aquarium and is not part of an established population. However, additional monitoring of Chicago harbors will continue to provide critical early warning.

Kolar and Lodge (2002) recently developed quantitative models of invasability characteristics of fishes from the Ponto-Caspian region. Based on their models, other fishes that could rise to pest status if they do establish in the Great Lakes include tyulka, Eurasian minnow, Black Sea silverside, European perch, and monkey goby.

Direct and Indirect Impacts

Invasive species affect Lake Michigan fishes both directly and indirectly. We illustrate direct impacts using round goby as an example and indirect impacts using dreissenid mussels, because few effects of the other species are known.


Round gobies can exert both positive and negative impacts on the nearshore fish community. They serve as prey for nearshore fishes, such as yellow perch (Truemper and Lauer 2005) and smallmouth bass, whose growth rates at age 0 have increased in Lake Erie since round gobies became abundant (Steinhart et al. 2004). Similarly, lake whitefish, lake trout, burbot, brown

trout, and coho salmon have been reported to consume round gobies. Although consumption of round gobies could be positive for predators, negative impacts of consumption of round gobies are also likely. Specifically, because round gobies >50 mm in length consume primarily dreissenid mussels, biomagnification of toxic substances, such as polychlorinated biphenyls and polychlorinated naphthalenes, through the food web is likely (Hanari et al. 2004).

Round gobies have essentially eliminated important nearshore benthic fishes, such as mottled sculpin and johnny darter (Janssen and Jude 2001; Lauer et al. 2004). The mechanisms behind this displacement include agonistic interactions for refuges (Dubs and Corkum 1996) and egg predation by round gobies on mottled sculpin nests (Janssen and Jude 2001). Because round gobies are voracious egg predators, they may negatively affect restoration efforts for native fishes by eating their eggs and fry (e.g., Chotkowski and Marsden 1999). For example, sturgeon eggs are consumed by round gobies (J. Nichols, unpublished data). Furthermore, round gobies are both more numerous and more-effective lake trout egg predators than native predators, such as crayfish (*Orconectes* spp.) and mottled sculpin (Fitzsimons et al. 2003).

Because dreissenid mussels filter phytoplankton from the water column, they compete directly with zooplankton for food (Fig. 2). Since dreissenid mussels invaded Lake Michigan, zooplankton densities, when first-feeding of yellow perch larvae occurs, have declined by an order of magnitude (Dettmers et al. 2003), indirectly resulting in reduced numbers of age-0 yellow perch in the fall (Clapp and Dettmers 2004).

Fig. 2. Simplified food-web structure of Lake Michigan in 1900 (a) and 2004 (b). Arrows represent the relative energy flow from one group to another, with thicker arrows representing greater energy flow. Dashed lines represent the reduced and declining contribution of *Diporeia* spp. to the trophic level immediately above. Species in **bold** text are invaders or intentionally introduced species that have established themselves in the food web.



The recent decline in *Diporeia* spp. (hereafter, diporeia as a common name) populations in southern Lake Michigan is another apparent indirect effect of dreissenid mussels. Although no specific mechanism for the decline has been demonstrated, diporeia populations did not start declining until zebra mussels were first detected (1989), and, by 1993, diporeia populations declined by over an order of magnitude in parts of the lake (Nalepa et al. 1998). This decline is relevant to the health of nearshore fishes because diporeia is an energy-rich food source and an important prey for several fishes, including alewife, yellow perch, and slimy sculpin (Wells 1980). Alewife condition declined by about 14% after 1995 (Madenjian et al. 2003), suggesting a link between reduced alewife condition and diporeia abundance. More recently, Madenjian et al. (2006) (also see the Status of Planktivore Populations chapter) reexamined the energy density of alewife using bomb calorimetry. Peak energy density of alewife >150 mm in length in the fall declined by 21%, from 9,641 J/g in 1979-1981 (Stewart and Binkowski 1986) to about 7,680 J/g in 2002-2003. In contrast, peak energy density of juvenile alewives, which do not consume diporeia, has remained at about 4,600 J/g. Salmonines must now consume greater numbers of large alewife than they did 30 years ago to achieve the same growth. This constraint may have important repercussions for the entire ecosystem as fishery managers seek optimal levels of predation by salmonines on alewives.

Numerous invasive species have established in Lake Michigan during the last 20 years (Fig. 2), and some of them exert strong impacts on the food web (Jude et al. 2005b). Although some invaders like the spiny water flea have limited discernable impacts, and others like the round goby can have positive impacts, the majority of impacts of invaders appears to be negative. The mechanisms for negative interactions are direct predation, as on lake trout eggs, or competition, as with other nearshore benthic fishes. Indirect interactions of dreissenid mussels, mediated through the food web, have probably negatively affected the recruitment of yellow perch and the condition of alewives. These indirect interactions have the potential for far-ranging impacts on the entire fish community (Fig. 2). Furthermore, after an invader has established itself, fishery managers are largely at the mercy of community interactions to determine if an invader will be a pest. Fishery managers face a very difficult task trying to manage fisheries that are part of an ever-changing set of food-web linkages mediated by invasive species.

Because there is great concern among fishery managers throughout the basin about the potential for new invaders to disrupt the existing community structure, an important emerging need will be tools to (1) predict which species are likely to invade, (2) predict which species are likely to become pests, and (3) especially prevent establishment of those likely to become pests. Without effective prevention measures, including but not limited to (1) eliminating transport by ballast water, (2) preventing planned introductions to other parts of the country that then spread through connecting waterways, and 3) establishing effective rapid-response plans for areas, such as ports and connecting channels, that are potential entry points, the fish community will remain at risk of disruption by invasive species.

THE STATE OF LAKE MICHIGAN IN 2005



SPECIAL PUBLICATION 08-02

THE STATE OF LAKE MICHIGAN IN 2005

Edited by

David F. Clapp

Charlevoix Great Lakes Station
96 Grant Street
Charlevoix, MI, 49720, U.S.A.

and

William Horns

Wisconsin Department of Natural Resources
101 S. Webster Street
Madison, WI, 53707-7921, U.S.A.

Citation (entire volume): Clapp, D.F., and W. Horns [EDS.]. 2008. The state of Lake Michigan in 2005. Great Lakes Fish. Comm. Spec. Pub. 08-02.


Citation (individual chapter): Wright, G., Honeyfield, D.C., and Faisal, M. 2008. Fish health. *In* The state of Lake Michigan in 2005. *Edited by* D.F. Clapp and W. Horns. Great Lakes Fish. Comm. Spec. Pub. 08-02. pp. 59-64.

Great Lakes Fishery Commission
2100 Commonwealth Blvd., Suite 100
Ann Arbor, MI 48105-1563

October 2008

ISSN 1090-1051

Printed on recycled paper.
SP08-02/10-2008/900



Lake Michigan depicting locations not otherwise identified in this publication. The lake basin is in grey, and treaty-ceded waters are depicted by diagonal lines.

LITERATURE CITED

- Adams, S.M. 1999. Ecological role of lipids in the health and success of fish populations. *In* Lipids in freshwater ecosystems. *Edited by* M.T. Arts and B.C. Wainman. Springer-Verlag, New York, NY. pp. 132-160.
- Albert, D.A., and Minc, L.D. 2001. Abiotic and floristic characterization of Laurentian Great Lakes' coastal wetlands. *Verh. Int. Ver. Limnol.* **27**(6): 3413-3419.
- Allen, P.J. 2000. A computer simulation model for the yellow perch population in the Indiana waters of Lake Michigan. M.Sc. thesis. Ball State University, Muncie, IN.
- Allen, P.J., and Lauer, T.E. 2005. Preliminary results of 2005 Ball State University yellow perch research in Indiana waters of Lake Michigan. Report of Ball State Univ., Div. Fish and Wildl. and Indiana Dept. Nat. Resour.
- Allen, P.J., Lauer, T.E., and McComish, T.S. 2005. Dynamics and models of the yellow perch in Indiana waters of Lake Michigan and near-shore fish community characteristics. Progress report for 2004-2005. Federal Aid Project F-18-R, Study 11, Year 3. Indiana Dept. Nat. Resour.
- Auer, N.A., Lepara, F.A., and Baker E. 2004. Assessment of remnant lake sturgeon populations in five Lake Michigan tributaries 2002-2003, with emphasis on physical river characteristics. Final report to the Great Lakes Fishery Trust. Project No. 2001.113. Mich. Technological Univ., Dept. Biol. Sci., Houghton, MI.
- Bartron, M.L., and Scribner, K.T. 2004. Temporal comparisons of genetic diversity in Lake Michigan steelhead, *Oncorhynchus mykiss*, populations: effects of hatchery supplementation. *Environ. Biol. Fishes* **69**: 395-407.
- Bassett, C. 1981. Management plan for lake sturgeon (*Acipenser fulvescens*) in the Indian River and Indian Lake, Alger and Schoolcraft Counties, Michigan. U.S. Forest Service, Manistique, MI.
- Bednarik, A.L. 2001. Undamming rivers: a review of the ecological impacts of dam removal. *Environ. Manage.* **27**(6): 803-814.
- Bence, J.R., and Ebener, M.P. 2002. Executive summary. *In* Summary status of lake trout and lake whitefish populations in the 1836 treaty-ceded waters of Lakes Superior, Huron and Michigan in 2000, with recommended yield and effort levels for 2001. *Edited by* J.R. Bence and M. Ebener. Technical Fisheries Committee, 1836 treaty-ceded waters of Lakes Superior, Huron, and Michigan.
- Benjamin, D.M., and Bence, J.R. 2003. Statistical catch-at-age framework for Chinook salmon in Lake Michigan, 1985-1996. Mich. Dept. Nat. Resour., Fish. Div. Res. Rep. 2066.

- Benoit, H.P., Johannsson, O.E., Warner, D.M., Sprules, W.G., and Rudstam, L.G. 2002. Assessing the impact of a recent predatory invader: The population dynamics, vertical distribution, and potential prey of *Cercopagis pengoi* in Lake Ontario. *Limnol. Oceanogr.* **47**: 626-635.
- Benson, A.C. 2004. Characterization of early life-history stages of lake sturgeon in the lower Peshtigo River, Wisconsin, and nearshore waters of Green Bay. M.Sc. thesis. Purdue Univ., West Lafayette, IN.
- Berejikian, B.A., Tezak, E.P., Schroder, S.L., Flagg, T.A., and Knudsen, C.M. 1999. Competitive differences between newly emerged offspring of captive-reared and wild coho salmon. *Trans. Am. Fish. Soc.* **128**(5): 832-839.
- Berejikian, B.A., Tezak, E.P., and Schroder, S.L. 2001. Reproductive behavior and breeding success of captively reared Chinook salmon. *North Am. J. Fish. Manage.* **21**(1): 255-260.
- Bertram, P., Forst, C., and Horvatin P. 2005. Developing indicators of ecosystem health. *In* State of Lake Michigan: ecology, health and Management. 2005. Ecovision World Mono. Ser., Aquatic Ecosystem Health Manage. Soc. pp. 505-519.
- Borgeson, D.P. 1970. Coho salmon status report 1967-1968. Michigan Dept. Nat. Resour., Fish. Div. Fish. Manage. Rep. No. 3.
- Brandt, S.B., Mason, D.M., MacNeill, D.B., Coates, T., and Gannon, J.E. 1987. Predation by alewives on larvae of yellow perch in Lake Ontario. *Trans. Am. Fish. Soc.* **116**: 641-645.
- Brandt, S.B., Mason, D.M., and Patrick, E.V. 1992. Spatially-explicit models of fish growth rate. *Fisheries* **17**: 23-33.
- Brazner, J.C. 1997. Regional, habitat and human development influences on coastal wetland and beach fish assemblages in Green Bay, Lake Michigan. *J. Great Lakes Res.* **23**(1): 36-51.
- Brazner, J.C., Campana, S.E., Tanner, D.K., and Schram, S.T. 2004. Reconstructing habitat use and wetland nursery origin of yellow perch from Lake Superior using otolith elemental analysis. *J. Great Lakes Res.* **30**(4): 492-507.
- Bremigan, M.T., Dettmers, J.M., and Mahan, A.L. 2003. Zooplankton selectivity by larval yellow perch in Green Bay, Lake Michigan. *J. Great Lakes Res.* **29**: 501-510.
- Brent, R.N., and Warren, G.J. 2005. Atrazine in the Lake Michigan ecosystem: monitoring results from the Lake Michigan mass balance study. *In* State of Lake Michigan: ecology, health and management. Ecovision World Mono. Ser., Aquatic Ecosystem Health Manage. Soc. pp. 157-179.

- Bronte, C.R., Holey, M.E., Breidert, B., Claramunt, R.M., Ebener, M.P., Fleischer, G.W., Hess, R., Martell, A., Olsen, E.J., McKee, P., and Toney, M.L. 2003a. Success of current strategies to re-colonize lake trout spawning reefs in northern Lake Michigan. Final Report, Project 1999.6, Great Lakes Fishery Trust.
- Bronte, C.R., Jonas, J.L., Holey, M.E., Eshenroder, R.L., Toney, M.L., McKee, P., Breidert, B., Claramunt, R.M., Ebener, M.P., Krueger, C.C., Wright, G., and Hess, R. 2003b. Possible impediments to lake trout restoration in Lake Michigan [online]. Available from <http://www.glfc.org/lakecom/lmc/lrestore.pdf> [accessed 12 September 2008].
- Bronte, C.R., Holey, M.E., Madenjian, C.P., Jonas, J.L., Claramunt, R.M., McKee, P.C., Toney, M.L., Ebener, M.P., Breidert, B., Fleischer, G.W., Hess, R., Martell, Jr., A.W., and Olsen, E.J. 2007. Relative abundance, site fidelity, and survival of adult lake trout in Lake Michigan from 1999-2001: implications for future restoration strategies. *North Am. J. Fish. Manage.* **27**(1): 137-155.
- Bronte, C.R., Krueger, C.C., Holey, M.E., Toney, M.L., Eshenroder, R.L., and Jonas, J.L. 2008. A guide for the rehabilitation of lake trout in Lake Michigan. *Great Lakes Fish. Comm. Misc. Pub.* 2008-01.
- Brown, R.W., Ebener, M., and Gorenflo, T. 1999. Great Lakes commercial fisheries: historical overview and prognosis for the future. *In* Great Lakes fisheries policy and management, a binational perspective. *Edited by* W.W. Taylor and C.P. Ferreri. Mich. State Univ. Press, East Lansing, MI. pp. 259-354.
- Brown, S.B., and Honeyfield, D.C. 2004. Fourth early mortality syndrome workshop research report [online]. Available from <http://www.glfc.org/research/reports/BrownEMS2004.pdf> [accessed 12 September 2008].
- Brown, S.B., Arts, M.T., Brown, L.R., Brown, M., Moore, K., Villella, M., Fitzsimons, J.D., Honeyfield, D.C., Tillitt, D.E., Zajicek, J.L., Wolgamood, M., and Hnath, J.G. 2005a. Can diet-dependent factors help explain fish-to-fish variation in thiamine-dependent early mortality syndrome? *J. Aquat. Anim. Health.* **17**: 36-47.
- Brown, S.B., Fitzsimons, J.D., Honeyfield, D.C., and Tillitt, D.E. 2005b. Implications of thiamine deficiency in Great Lake salmonines. *J. Aquat. Anim. Health.* **17**: 113-124.
- Brown, S.B., Honeyfield, D.C., Hnath, J., Wolgamood, M., Marquenski, S.V., Fitzsimons, J.D., and Tillitt, D.E. 2005c. Thiamine status in adult salmonines in the Great Lakes. *J. Aquat. Anim. Health.* **17**: 59-64.
- Bruch, R.M. 1999. Management of lake sturgeon on the Winnebago system: long-term impacts of harvest and regulations on population structure. *J. Appl. Ichthyol.* **15**: 142-152.

- Carl, L.M. 1982. Natural reproduction of coho salmon and Chinook salmon in some Michigan streams. *North Am. J. Fish. Manage.* **4**: 375-380.
- Carl, L.M. 1983. Density, growth, and change in density of coho salmon and rainbow trout in three Lake Michigan tributaries. *Can. J. Zool.* **61**: 1120-1127.
- Charlebois, P.M., Marsden, J.E., Goettel, R.G., Wolfe, R.K., Jude, D.J., and Rudnicka, S. 1997. The round goby, *Neogobius melanostomus* (Pallas), a review of European and North American literature. Illinois-Indiana Sea Grant Program and Illinois Natural History Survey. INHS Special Pub. No. 20.
- Charlebois, P.M., Raffenberg, M.J., and Dettmers, J.M. 2001. First occurrence of *Cercopagis pengoi* in Lake Michigan. *J. Great Lakes Res.* **27**: 258-261.
- Chiotti, J.A. 2004. Evaluation of spawning habitat, juvenile habitat, and larval drift of lake sturgeon (*Acipenser fulvescens*) in the Big Mansistee River, Michigan. M.Sc. thesis, Mich. Tech. Univ., Houghton, MI.
- Chotkowski, M.A., and Marsden, J.E. 1999. Round goby predation on lake trout eggs and fry: field predictions from laboratory experiments. *J. Great Lakes Res.* **25**: 26-35.
- Christie, W.J., Collins, J.J., Eck, G.W., Goddard, C.I., Hoenig, J.M., Holey, M., Jacobson, L.D., MacCallum, W., Nepszy, S. J., O'Gorman, R., and Selgeby, J. 1987. Meeting future information needs for Great Lakes fisheries management. *Can. J. Fish. Aquat. Sci.* **44**(Suppl. 2): 439-447.
- Christie, G.C., and Goddard, C.L. 2003. Sea lamprey symposium (SLIS II): advances in the integrated management of sea lamprey in the Great Lakes. *J. Great Lakes Res.* **29**(Suppl. 1): 1-14.
- Clapp, D.F., and Dettmers, J.M. 2004. Yellow perch research and management in Lake Michigan: evaluating progress in a collaborative effort, 1997-2001. *Fisheries* **29**(11): 11-19.
- Clapp, D.F., Schneeberger, P.J., Jude, D.J., Madison, G., and Pistis, C. 2001. Monitoring round goby (*Neogobius melanostomus*) population expansion in eastern and northern Lake Michigan. *J. Great Lakes Res.* **27**: 335-341.
- Clapp, D.F., Schneeberger, P.J., O'Neal, R.P., Lychwick, T.J., Belonger, B., and Shroyer, S.M. 2005. Inshore fish community. *In* The state of Lake Michigan in 2000. *Edited by* M.E. Holey and T.N. Trudeau. Great Lake Fish. Comm. Spec. Pub. 05-01, pp. 49-58.
- Claramunt, R.M., Breidert, B., Clapp, D.F., Elliott, R.F., Madenjian, C.P., Peeters, P., Robillard, S., and Wright, G. 2004. Status of Chinook salmon in Lake Michigan, 1985-2003 [online]. Available from http://www.michigandnr.com/publications/pdfs/Fishing/research/Charlevoix/132_Status-of-CHS-Report_2007_final.pdf [accessed 15 September 2008].

- Claramunt, R.M., Jonas, J.L., Fitzsimons, J.D., and Marsden, J.E. 2005. Influences of spawning habitat characteristics and interstitial predators on lake trout egg deposition and mortality. *Trans. Am. Fish. Soc.* **134**:1048-1057.
- Claramunt, R.M., Breidert, B., Clapp, D.F., Elliott, R.F., Madenjian, C.P., Peeters, P., Robillard, S., Warner, D.M., and Wright, G. 2006. Status of Chinook salmon in Lake Michigan, 1985-2005 [online]. Available from http://www.michigandnr.com/publications/pdfs/Fishing/research/Charlevoix/122_SWG_Report_2006.pdf [accessed 15 September 2008].
- Coakley, J.P., Rasul, N., Ioannou, S.E., and Brown, G.E. 2002. Soft sediment as a constraint on the spread of the zebra mussel in western Lake Erie: processes and impacts. *Aquat. Ecosyst. Health Manage.* **5**: 329-343.
- Coble, D.W., Brueswitz, R.E., Fratt, T.W., and Scheirer, J.W. 1990. Lake trout, sea lamprey, and overfishing in the upper Great Lakes: a review and reanalysis. *Trans. Am. Fish. Soc.* **119**: 985-995.
- Cochrane, G.R., and Lafferty, K.D. 2002. Use of acoustic classification of sidescan sonar data for mapping benthic habitat in the Northern Channel Islands, California. *Coastal Shelf Research* **22**: 683-690.
- Coon, T.G. 1999. Ichthyofauna of the Great Lakes basin. *In* Great Lakes fisheries policy and management a binational perspective. *Edited by* W. W. Taylor and C. P. Ferreri. Mich. State Univ. Press, East Lansing, MI. pp. 55-72.
- Council of Great Lakes Governors. 1985. The Great Lakes Charter [online]. Available from <http://www.cglg.org/projects/water/docs/GreatLakesCharter.pdf> [accessed 9 September 2008].
- Council of Great Lakes Governors. 2001. The Great Lakes Charter Annex [online]. Available from <http://www.deq.state.mi.us/documents/deq-ogl-Annex2001.pdf> [accessed 9 September 2008].
- Creque, S.M. 2002. Using landscape-scale habitat models to predict potential abundance of potamodromous fishes above dams on Great Lakes tributaries. M.Sc. thesis, Univ. Mich., Ann Arbor, MI.
- Daugherty, D.J., and Sutton, T.M. 2004. Assessment of remnant lake sturgeon populations in the St. Joseph and Kalamazoo Rivers. Final report to the Great Lakes Fishery Trust. Project No. 2001.113. Purdue Univ., Dept. Forestry Nat. Resour., West Lafayette, IN.
- DeHaan, P.W. 2003. Demographic and life history characteristics of remnant lake sturgeon populations in the upper Great Lakes basin: inferences based on genetic analysis. M.Sc. thesis, Mich. State Univ., East Lansing, MI.

- Dettmers, J.M., Raffenberg, M.J., and Weis, A.K. 2003. Exploring zooplankton changes in southern Lake Michigan: implications for yellow perch recruitment. *J. Great Lakes Res.* **29**: 355-364.
- Dexter, J.L., and LeDet, N.D. 1997. Estimates of fish passage on the St. Joseph River in 1993 using time-lapse video recording. Michigan Dept. Nat. Resour., Fish. Div. Spec. Rep. No. 23.
- Dubs, D.O.L., and Corkum, L.D. 1996. Behavioral interactions between round gobies (*Neogobius melanostomus*) and mottled sculpins (*Cottus bairdi*). *J. Great Lakes Res.* **22**: 838-844.
- Dufour, E., Patterson, W.P., Höök, T.O., and Rutherford, E.S. 2005. Early life history of Lake Michigan alewives (*Alosa pseudoharengus*) inferred from intra-otolith stable isotope ratios. *Can. J. Fish. Aquat. Sci.* **62**(10): 2362-2370.
- Ebener, M.P., and Copes, F.A. 1985. Population statistics, yield estimates, and management considerations for two lake whitefish stocks in Lake Michigan. *North Am. J. Fish. Manage.* **5**: 435-448.
- Ebener, M.P., Bence, J.R., Newman, K., and Schneeberger, P.J. 2005. Application of statistical catch-at-age models to assess lake whitefish stocks in the 1836 treaty-ceded waters of the upper Great Lakes. *In* Proceedings of a workshop on the dynamics of lake whitefish (*Coregonus clupeaformis*) and the amphipod *Diporeia* spp. in the Great Lakes. *Edited by* L.C. Mohr and T.F. Nalepa, Great Lakes Fish. Comm. Tech. Rep. No. 66. pp. 271-309.
- Eissa, A., Elsayed, E., and Faisal, M. 2004. First record of *Renibacterium salmoninarum* isolation from the sea lamprey (*Petromyzon marinus*). Abstract, Proceeding of the 29th Annual Eastern Fish Health Workshop, NC, March 2004.
- Elliott, R.F. 1993. Feeding habits of Chinook salmon in eastern Lake Michigan. M.Sc. thesis, Michigan State University, East Lansing, MI.
- Environmental Working Group. 2006. Toxic fire retardants (PBDEs) in human breast milk [online]. Available from <http://www.ewg.org/reports/mothersmilk/> [accessed 02 October 2008].
- Eshenroder, R.L., Holey, M.E., Gorenflo, T.K., and Clark, Jr., R.D. 1995. Fish-community objectives for Lake Michigan. *Great Lakes Fish. Comm. Spec. Pub.* 95-3.
- Evans, M.S. 1988. *Bythotrephes cederstroemi* its new appearance in Lake Michigan. *J. Great Lakes Res.* **14**: 234-240.

- Faisal, M., and Hnath, J.G. 2005. Fish health and diseases issues in the Laurentian Great Lakes. *In* Health And Diseases of Aquatic Organisms: Bilateral Perspectives. *Edited by* R.C. Cipriano, I.S. Shchelkunov, and M. Faisal. Proceedings of the Second Bilateral Conference between Russia and the United States. 21-28 September 2003. Shepherdstown, WV. Mich. State Univ. Press. East Lansing, MI. pp. 331-350.
- Falkner, G., and Falkner, R. 2000. Objectivistic views in biology: an obstacle to our understanding of self-organisation processes in aquatic ecosystems. *Freshwater Biol.* **44**(3): 553-559.
- Fitzsimons, J.D., Brown, S.B., Honeyfield, D.C., and Hnath, J.G. 1999. A review of early mortality syndrome (EMS) in Great Lakes salmonids: a relationship with thiamine deficiency. *Ambio* **28**: 9-15.
- Fitzsimons, J.D., Marsden, J.E., Ellrott, B.J., Jonas, J., and Claramunt, R.M. 2003. Effects of egg and fry predators on lake trout recruitment in Lake Michigan [online]. Available from www.glfc.org/research/rcr.php [accessed 9 September 2008].
- Fleischer, G.W., DeSorcie, T.J., and Holuszko, J.D. 2001. Lake-wide distribution of *Dreissena* in Lake Michigan. *J. Great Lakes Res.* **27**: 252-257.
- Fleischer, G.W., Madenjian, C.P., Elliott, R.F., and Toney, M.L. 2005. Planktivores. *In* The state of Lake Michigan in 2000. *Edited by* M.E. Holey and T.N. Trudeau. Great Lakes Fish. Comm. Spec. Pub. 05-01, pp. 16-20.
- Foley, J.A., DeFries, R., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coe, M.T., Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard, E.A., Kucharik, C.J., Monfreda, C., Patz, J.A., Prentice, I.C., Ramankutty, N., and Snyder, P.K.. 2005. Global consequences of land use. *Science* **309**(5734): 570-574.
- Francis, J.T., Robillard, S.R., and Marsden, J.E. 1996. Yellow perch management in Lake Michigan: a multi-jurisdictional challenge. *Fisheries* **21**(2): 18-20.
- Glover, D.C., Dettmers, J.M., Wahl, D.H., and Clapp, D.F. 2008. Yellow perch stock structure in Lake Michigan: an analysis using mark-recapture data. *Can. J. Fish. Aquat. Sci.* **65**: 1919-1930.
- Graeb, B.D.S., Dettmers, J.M., Wahl, D.H., and Cáceres, C.E. 2004. Fish size and prey availability affect growth, survival, prey selection, and foraging behavior of larval yellow perch. *Trans. Am. Fish. Soc.* **133**: 504-514.
- Great Lakes Basin Water Resources Compact. 2005. Great Lakes—St. Lawrence River basin water resources compact [online]. Available from <http://www.deq.state.mi.us/documents/deq-ogl-Annex2001-Compact-12-13-05.pdf> [accessed 9 September 2008].

- Great Lakes Fishery Commission. 1997. A Joint Strategic Plan for Management of Great Lakes Fisheries [online]. Available from <http://www.glfc.org/fishmgmt/jsp97.pdf> [accessed 9 September 2008].
- Great Lakes Fishery Commission. 2005. Lake Michigan GIS [online]. Available from http://www.glfc.org/glgis/support_docs/html/lake_GISs/LMGIS_index.htm [accessed 9 September 2008].
- Great Lakes Regional Collaboration. 2005. Great Lakes Regional Strategy [online]. Available from <http://www.glrc.us/> [accessed 9 September 2008].
- Grigorovich, I.A., Colautti, R.I., Mills, E.L., Holeck, K., Ballert, A.G., and MacIsaac, H.J. 2003. Ballast-mediated animal introductions in the Laurentian Great Lakes: retrospective and prospective analyses. *Can. J. Fish. Aquat. Sci.* **60**: 740-756.
- Gunderman, B.J. 2001. Population dynamics of lake sturgeon (*Acipenser fulvescens* Rafinesque) in the Manistee River. M.Sc. thesis, Central Mich. Univ., Mount Pleasant, MI.
- Gunderman, B., and Elliott, R. 2004. Assessment of remnant lake sturgeon populations in the Green Bay basin, 2002-2003. Final report to the Great Lakes Fishery Trust. Project Number 2001.113. U.S. Fish Wildl. Serv., Green Bay Fish. Resour. Office, New Franken, WI.
- Gunderson, J.L., Klepinger, M.R., Bronte, C.R., and Marsden, J.E. 1998. Overview of the international symposium on Eurasian ruffe (*Gymnocephalus cernuus*) biology, impacts, and control. *J. Great Lakes Res.* **24**: 165-169.
- Hanari, N., Kannan, K., Horii, Y., Taniyasu, S., Yamashita, N., Jude, D.J., and Berg, M.B. 2004. Polychlorinated naphthalenes and polychlorinated biphenyls in benthic organisms of a Great Lakes food chain. *Arch. Environ. Contam. Toxicol.* **47**: 84-93.
- Hansen, M.J. 1999. Lake trout in the Great Lakes: basinwide stock collapse and binational restoration. *In* Great Lakes fisheries policy and management, a binational perspective. *Edited by* W.W. Taylor and C.P. Ferreri. Mich. State Univ. Press, East Lansing, MI. pp. 417-454.
- Hansen, M.J., and Holey, M.E. 2002. Ecological factors affecting sustainability of Chinook and coho salmon populations in the Great Lakes. *In* Sustaining North American salmon: perspectives across regions and disciplines. *Edited by* K.D. Lynch, M.L. Jones, and W.W. Taylor. Am. Fish. Soc. pp. 155-179.
- Hanson, D. 2005. Lake Michigan recreational fishery creel database 2004 summary report. Minutes of the 2005 meeting of the Lake Michigan Committee, Great Lakes Fishery Commission, Ann Arbor, MI.

- Hart, D.D., Johnson, T.E., Bushaw-Newton K.L., Horwitz, R.J., Bednarek, A.T., Charles, D.F., Kreeger, D.A., and Velinsky, D.J. 2002. Dam removal: challenges and opportunities for ecological research and river restoration. *Bioscience* **52**(8): 669-681.
- Hay-Chmielewski, L., and Whelan, G. [EDS.]. 1997. Lake sturgeon rehabilitation strategy. Mich. Dept. Nat. Resour., Fisheries Div., Spec. Rep. 18, Lansing, MI.
- Hecky, R.E., Smith, R.E.H., Barton, D.R., Guildford, S.J., Taylor, W.D., Charlton, M.N., and Howell, T. 2004. The nearshore phosphorus shunt: a consequence of ecosystem engineering by dreissenids in the Laurentian Great Lakes. *Can. J. Fish. Aquat. Sci.* **61**: 1285-1293.
- Heins, D.C., and Baker, J.A. 2003. Reduction of egg size in natural populations of three spine stickleback infected with a cestode macroparasite. *J. Parasitol.* **89**: 1-6.
- Hensler, S. 2004. Yellow perch recruitment, zooplankton changes, and larval fish catchability comparisons in eastern Lake Michigan. M.Sc. thesis, Univ. Mich., Ann Arbor, MI.
- Hesse, J.A. 1994. Contributions of hatchery and natural Chinook salmon to the eastern Lake Michigan fishery, 1992-93. Mich. Dept. Nat. Resour., Fish. Div., Fish. Res. Rep. No 2013.
- Heyer, C.J., Miller, T.J., Binkowski, F.P., Caldarone, E.M., and Rice, J.A. 2001. Maternal effects as a recruitment mechanism in Lake Michigan yellow perch (*Perca flavescens*). *Can. J. Fish. Aquat. Sci.* **58**: 1477-1487.
- Hirethota, P.S., Burzinski, T.E., and Eggold, B.T. 2005. Changing habitat and biodiversity of the lower Milwaukee River and estuary. Wisc. Dept. Nat. Resour. PUB-FIT-511-2005. Milwaukee, WI.
- Hogler, S., and Surendonk, S. 2004. Return, size, and age of steelhead at the Besadny Anadromous Fisheries Facility, 2004 [online]. Available from http://dnr.wi.gov/fish/lakemich/BAFF_2004.pdf [accessed 24 September 2008].
- Holey, M.E., Elliott, R.F., Marquenski, S.V., Hnath, J.G., and Smith, K.D. 1998. Chinook salmon epizootics in Lake Michigan: possible contributing factors and management implications. *J. Aquat. Anim. Health* **10**: 202-210.
- Holey, M.E., Rybicki, R.R., Eck, G.W., Brown, Jr., E.H., Marsden, J.E., Lavis, D.S., Toneys, M.L., Trudeau, T.N., and Horrall, R.M. 1995. Progress toward lake trout restoration in Lake Michigan. *J. Great Lakes Res.* **21**(Suppl. 1): 128-151.
- Holey, M.E., and Trudeau, T.N. [EDS.]. 2005. The state of Lake Michigan in 2000. *Great Lakes Fish. Comm. Spec. Pub.* 05-01.

- Honeyfield, D.C., Hinterkopf, J.P., Fitzsimons, J.D., Brown, S.B., Tillitt, D.E., and Zajicek, J. 2005. Development of thiamine deficiencies and early mortality syndrome (EMS) in lake trout by feeding experimental and feral fish diets containing thiaminase. *J. Aquat. Anim. Health*. **17**:4-12.
- International Joint Commission. 1988. Revised Great Lakes Water Quality Agreement of 1978, as amended by protocol signed November 18, 1987/September 1989 [online]. Available from <http://www.ijc.org/rel/agree/quality.html> [accessed 9 September 2008].
- Janssen, J., and Jude, D.J. 2001. Recruitment failure of mottled sculpin *Cottus bairdi* in southern Lake Michigan induced by the newly introduced round goby, *Neogobius melanostomus*. *J. Great Lakes Res.* **27**: 319-328.
- Janssen, J., Jude, D.J., Edsall, T.A., Paddock, R.W., Wattrus, N., Toneys, M., and McKee, P. 2006. Evidence of lake trout reproduction at Lake Michigan's mid-lake reef complex. *J. Great Lakes Res.* **32**(4): 749-763.
- Johnson, D.A., Weisser, J.W., and Bills, T.D. 1999. Sensitivity of lake sturgeon (*Acipenser fulvescens*) to the lampricide 3-trifluoromethyl-4-nitrophenol (TFM) in field and laboratory exposures. *Great Lakes Fish. Comm., Tech. Rep.* 62.
- Johnson, D.C., and Hnath, J.G. 1991. Lake Michigan Chinook salmon mortality-1988. Mich. Dept. Nat. Resour., Fish. Div., Tech. Rep. No. 91-4.
- Jonas, J.L., Clapp, D.F., Bence, J.R., and Holey, M.E. 2005a. Salmonine community. *In* The state of Lake Michigan in 2000. *Edited* by M.E. Holey and T.N. Trudeau. *Great Lakes Fish. Comm. Spec. Pub.* 05-01, pp. 33-47.
- Jonas, J.L., Claramunt, R.M., Fitzsimons, J.D., Marsden, J.E., and Elliott, B.J. 2005b. Estimates of egg deposition and effects of lake trout (*Salvelinus namaycush*) egg predators in three regions of the Great Lakes. *Can. J. Fish. Aquat. Sci.* **62**(10): 2254-2264.
- Jonas, J.L., Schneeberger, P.J., Clapp, D.F., Wolgamood, M., Wright, G., and Lasee, B. 2002. Presence of the BKD-causing bacterium *Renibacterium salmoninarum* in lake whitefish and bloaters in the Laurentian Great Lakes. *Arch. Hydrobiol., Spec. Issues, Advanc. Limnol.* **57**: 447-452.
- Jude, D.J. 2001. Round and tubenose gobies: 10 years with the latest Great Lakes phantom menace. *Dreissena* **11**(4): 1-14.
- Jude, D.J., Albert, D., Uzarski, D.G. and Brazner, J. 2005a. Lake Michigan's coastal wetlands: distribution, biological components with emphasis on fish, and threats. *In* State of Lake Michigan: ecology, health and management. *Edited* by T. Edsall and M. Munawar. *Ecovision World Mono. Ser., Aquatic Ecosystem Health Manage. Soc.* pp. 439-477.

- Jude, D.J., Janssen, J., and Stoermer, E. 2005b. The uncoupling of trophic food webs by invasive species in Lake Michigan. *In* The state of Lake Michigan: ecology, health, and management. *Edited by* T. Edsall and M. Munawar. Ecovision World Mono. Ser., Aquatic Ecosystem Health Manage. Soc. pp. 311-348.
- Keller, M., Smith, K.D., and Rybicki, R.W. 1990. Review of salmon and trout management in Lake Michigan. Mich. Dept. Nat. Resour., Fish. Div., Fish. Spec. Rep. No 14.
- King, Jr., E.L. 1980. Classification of sea lamprey (*Petromyzon marinus*) attack marks on Great Lakes lake trout (*Salvelinus namaycush*). Can. J. Fish. Aquat. Sci. **37**: 1989-2006.
- Kocik, J. F., and Jones, M. L. 1999. Pacific salmonines in the Great Lakes basin. *In* Great Lakes fisheries policy and management: a binational perspective. *Edited by* W.W. Taylor and C.P. Ferreri. Mich. State Univ. Press, East Lansing, MI. pp. 455-488.
- Kolar, C.S., and Lodge, D.M. 2002. Ecological predictions and risk assessment for alien fishes in North America. Science **298**: 1233-1236.
- Kostow, K.E., Marshall, A.R., and Phelps, S.R. 2003. Naturally spawning hatchery steelhead contribute to smolt production but experience low reproductive success. Trans. Am. Fish. Soc. **132**(4): 780-790.
- Krueger, C.C., Perkins, D.L., Mills, E.L. and Marsden, J.E. 1995. Predation by alewives on lake trout fry in Lake Ontario: role of an exotic species in preventing restoration of a native species. J. Great Lakes Res. **21**(Supple. 1): 458-469.
- Lake Michigan Lake Trout Technical Committee. 1985. A draft lakewide plan for lake trout restoration in Lake Michigan. *In* Minutes of the 1985 Annual Meeting of the Lake Michigan Committee. Great Lakes Fishery Commission, Ann Arbor, MI.
- Lallaman, J.J. 2003. Stock assessment and summer movement patterns of lake sturgeon (*Acipenser fulvescens*) in the Manistee River, MI. M.Sc. thesis, Central Mich. Univ., Mount Pleasant, MI.
- Lauer, T.E., Allen, P.J., and McComish, T.S. 2004. Changes in mottled sculpin and johnny darter trawl catches after the appearance of round gobies in the Indiana waters of Lake Michigan. Trans. Am. Fish. Soc. **133**:185-189.
- Lauer, T.E., Shroyer, S.M., Kilpatrick, J.M., McComish, T.S., and Allen, P.J. 2005. Yellow perch length-fecundity and length-egg size relationships in Indiana waters of Lake Michigan. North Am. J. Fish. Manage. **25**: 791-796.
- Lavis, D.S., Henson, M.P., Johnson, D.A., Koon, E.M., and Ollila, D.J. 2003. A case history of sea lamprey control in Lake Michigan: 1979-1999. J. Great Lakes Res. **29**(Suppl. 1): 584-598.

- Laxson, C.L., McPhedran, K.N., Makarewicz, J.C., Telesh, I.V. 2003. Effects of the non-indigenous cladoceran *Cercopagis pengoi* on the lower food web of Lake Ontario. *Freshwater Biol.* **48**: 2094-2106.
- Lehman, J.T. 1991. Causes and consequences of cladoceran dynamics in Lake Michigan: implications of species invasion by *Bythotrephes*. *J. Great Lakes Res.* **17**: 437-445.
- Lehman, J.T., and Cáceres, C.E. 1993. Food-web responses to species invasion by a predatory invertebrate: *Bythotrephes* in Lake Michigan. *Limnol. Oceanogr.* **38**: 879-891.
- Lessard, J.L. 2001. Temperature effects of dams on coldwater fish and macroinvertebrate communities in Michigan. Mich. Dept. Nat. Resour., Fish. Div., Fish. Res. Rep. No 2058.
- MacLean, D.G., and Yoder, W.G. 1970. Kidney disease among Michigan salmon in 1967. *Prog. Fish-Cult.* **32**: 26-30.
- Madenjian, C.P., Elliott, R.F., DeSorcie, T.J., Stedman, R.M., O'Connor, D.V., and Rottiers, D.V. 2000. Lipid concentrations in Lake Michigan fishes: seasonal, spatial, ontogenetic, and long-term trends. *J. Great Lakes Res.* **26**: 427-444.
- Madenjian, C.P., Fahnenstiel, G.L., Johengen, T.H., Nalepa, T.F., Vanderploeg, H.A., Fleischer, G.W., Schneeberger, P.J., Benjamin, D.M., Smith, E.B., Bence, J.R., Rutherford, E.S., Lavis, D.S., Robertson, D.M., Jude, D.J., and Ebener, M.P. 2002. Dynamics of the Lake Michigan food web, 1970-2000. *Can. J. Fish. Aquat. Sci.* **59**: 736-753.
- Madenjian, C.P., Holuszko J.D., and Desorcie, T.J. 2003. Growth and condition of alewives in Lake Michigan, 1984-2001. *Trans. Am. Fish. Soc.* **132**: 1104-1116.
- Madenjian, C., Bunnell, D.B., DeSorcie, T.J., Holuszko, J.D., and Adams, J.V. 2005a. Status of prey fish populations in Lake Michigan, 2004. *In* Minutes of the 2005 Annual Meeting of the Lake Michigan Committee. Great Lakes Fishery Commission, Ann Arbor, MI.
- Madenjian, C.P., Höök, T.O., Rutherford, E.S., Mason, D.M., Croley, II, T.E., Szalai, E.B. and Bence J.R. 2005b. Recruitment variability of alewives in Lake Michigan. *Trans. Am. Fish. Soc.* **134**: 218-230.
- Madenjian, C.P., Pothoven, S.A., Dettmers, J.M., and Holuszko, J.D.. 2006. Changes in seasonal energy dynamics of alewife (*Alosa pseudoharengus*) in Lake Michigan after invasion of dreissenid mussels. *Can. J. Fish. Aquat. Sci.* **63**: 901-902.

- Makauskas, D., and Allen, P.J. 2004. Status of yellow perch in Lake Michigan and Yellow Perch Task Group. *In* Minutes of the 2005 Annual Meeting of the Lake Michigan Committee. Great Lakes Fishery Commission, Ann Arbor, MI.
- Marsden, J. E. 1992. The zebra mussel invasion. *Aquaticus* **23**: 19-29.
- May, B., and Marsden, J.E. 1992. Genetic identification and implications of another invasive species of dreissenid mussel in the Great Lakes. *Can. J. Fish. Aquat. Sci.* **49**: 1501-1506.
- McCarty, H.B., Schofield, J., Miller, K., Brent, R.N., Van Hoof, P., and Eadie, B. 2004. Results of the Lake Michigan mass balance study: polychlorinated biphenyls and trans-nonachlor data report. EPA 905 R-01-011, Chicago, IL.
- Miller, L.M. 2003. Microsatellite DNA loci reveal genetic structure of yellow perch in Lake Michigan. *Trans. Am. Fish. Soc.* **132**: 503-513.
- Mills, E.L., Leach, J.H., Carlton, J.T., and Secor, C.L. 1993. Exotic species in the Great Lakes: a history of biotic crises and anthropogenic introductions. *J. Great Lakes Res.* **19**: 1-54.
- Minns, C.K., Randall, R.G., Moore, J.E., and Cairns, V.W. 1996. A model simulating the impact of habitat supply limits on northern pike, *Esox lucius*, in Hamilton Harbour, Lake Ontario. *Can. J. Fish. Aquat. Sci.* **53**(Suppl. 1): 20-34.
- Mistak, J.L., Hayes, D.B., and Bremigan, M.T. 2003. Food habits of salmonines above and below Stronach Dam in the Pine River, Michigan. *Environ. Biol. Fishes* **67**(2): 179-190.
- Mohr, L.C., and Ebener, M.P. 2005. Status of lake whitefish (*Coregonus clupeaformis*) in Lake Huron. *In* Proceedings of a workshop on the dynamics of lake whitefish (*Coregonus clupeaformis*) and the amphipod *Diporeia* spp. in the Great Lakes. *Edited by* L.C. Mohr and T.F. Nalepa, Great Lakes Fish. Comm. Tech. Rep. No. 66. pp. 105-125.
- Mullett, K.M., Heinrich J.W., Adams, J.V., Young, R.J., Henson, M.P., McDonald, R.B., and Fodale, M.F. 2003. Estimating lake-wide abundance of spawning-phase sea lampreys (*Petromyzon marinus*) in the Great Lakes: extrapolating from sampled streams using regression models. *J. Great Lakes Res.* **29**(Suppl. 1): 240-252.
- Murphy, E. and Whittle, D.M. 2004. Environmental indicators of progress [online]. Available from http://binational.net/bns/2004/2004glbts_11a.pdf [accessed 9 September 2008].
- Nalepa, T.F., Hartson, D.J., Fanslow, D.L., Lang, G.A., and Lozano, S.L. 1998. Decline of benthic macroinvertebrate populations in southern Lake Michigan, 1980-1993. *Can. J. Fish. Aquat. Sci.* **55**: 2402-2413.

- Nalepa, T.F., Schloesser, D.W., Pothoven, S.A., Hondorp, D.W., Fanslow, D.L., Tuchman, M.L., and Fleischer, G.W. 2001. First finding of the amphipod *Echinogammarus ischnus* and the mussel *Dreissena bugensis* in Lake Michigan. *J. Great Lakes Res.* **27**: 384-391.
- Newcomb, T.J. 1998. Productive capacity of the Betsie River watershed for steelhead smolts. Ph.D. thesis. Mich. State Univ., East Lansing, MI.
- Ogle, D.H. 1998. A synopsis of the biology and life history of ruffe. *J. Great Lakes Res.* **24**: 170-185.
- O’Gorman, R., Gorman, O., and Bunnell, D. 2005. Great Lakes prey fish populations: a cross-basin view of status and trends in 2004. *In* Minutes of the 2005 Annual Meeting of the Common Session for the Upper Lakes. Great Lakes Fishery Commission, Ann Arbor, MI.
- Patriarche, M.H. 1980. Movement and harvest of coho salmon in Lake Michigan, 1978-1979. Mich. Dept. Nat. Resour., Fish. Div., Fish. Res. Rep. No. 1889.
- Peeters, P. and Royseck, K. 2003. Harvest, age, and size at age of Chinook and coho salmon at Strawberry Creek Weir and Besadny Anadromous Fisheries Facility, fall 2003 [online]. Available from http://dnr.wi.gov/fish/lakemich/SCW_BAFF_Report_Fall_2003.pdf [accessed 24 September 2008].
- Peterman, R.M. and Anderson, J.L. 1999. Decision analysis: a method for taking uncertainties into account in risk-based decision making. *Human Ecol. Risk Assess.* **5**: 231-244.
- Peters, A.K., Jones, M.L., Honeyfield, D., and Bence, J.R. 2007. Monitoring energetic status of Lake Michigan Chinook salmon using water content as a predictor of whole-fish lipid content. *J. Great Lakes Res.* **33**: 253-263.
- Peterson, D.L., Gunderman, B., and Vecsei, P. 2002. Lake sturgeon of the Manistee River: a current assessment of spawning stock size, age, and growth. *In* Biology, management, and protection of North American sturgeon. *Edited by* W. Van Winkle, P. Anders, D.H. Secor, and D. Dixon. *Am. Fish. Soc. Symp.* **28**: 175-182.
- Peterson, D.L., and Vecsei, P. 2004. Lake sturgeon of the Muskegon River: population dynamics and life history. Final report for the Great Lakes Fishery Trust. Project No. 2001.113. Univ. of Georgia, Warnell School of Forest Resources, Athens, GA.
- Poole, G.C., and Berman, C.A. 2001. An ecological perspective on instream temperature. Natural heat dynamics and the mechanisms of human caused thermal degradation. *Environ. Manage.* **27**(6): 787-802.

- Pothoven, S.A. 2005. Changes in lake whitefish diet in Lake Michigan, 1998-2001. *In* Proceedings of a workshop on the dynamics of lake whitefish (*Coregonus clupeaformis*) and the amphipod *Diporeia* spp. in the Great Lakes. *Edited by* L.C. Mohr and T.F. Nalepa, Great Lakes Fish. Comm. Tech. Rep. No. 66. pp. 127-140.
- Pothoven, S.A., Fahnenstiel, G.L., and Vanderploeg, H.A. 2003. Population characteristics of *Bythotrephes* in Lake Michigan. *J. Great Lakes Res.* **29**: 145-156.
- Pothoven, S.A., Nalepa, T.F., Schneeberger, P.J., Brandt, S.B. 2001. Changes in diet and body condition of lake whitefish in southern Lake Michigan associated with changes in benthos. *North Am. J. Fish. Manage.* **21**: 876-883.
- Raiffa, H. 1968. Decision analysis: introductory lectures on choices under uncertainty. Addison-Wesley, Don Mills, ON.
- Rand, P.S., Stewart, D.J., Seelbach, P.W., Jones, M.L., and Wedge, L.R. 1993. Modeling steelhead population energetics in Lakes Michigan and Ontario. *Trans. Am. Fish. Soc.* **122**: 977-1001.
- Rosenberg, A.A., and Restrepo, V.R. 1994. Uncertainty and risk evaluation in stock assessment advice for U.S. marine fisheries. *Can. J. Fish. Aquat. Sci.* **51**: 2715-2720.
- Rutherford, E.S., Marshall E., Clapp, D., Horns, W., Gorenflo, T., and Trudeau, T. 2004. Lake Michigan Environmental Objectives (draft report) [online]. Available from <http://www.glfc.org/research/reports/RutherfordEOs.pdf> [accessed 9 September 2008].
- Rutherford, E.S., Thorrold, S.R., Swank, D.R., Woldt, A.P., and Godby, J.R., N.G. 1999. Abundance, production and harvest of anadromous salmonids in the Au Sable, Manistee and Muskegon Rivers. Progress Report for # 231663, Amendment # 14, FY 96-97, Project 231663, Michigan Dept. Natural Resources, Fish. Div.
- Rybicki, R.W., and Clapp, D.F. 1996. Diet of Chinook salmon in eastern Lake Michigan, 1991-1993. *Mich. Dept. Nat. Resour., Fish. Div., Fish. Tech. Rep.* No. 2027.
- Scheerer, P.D., and Taylor, W.W. 1985. Population dynamics and stock differentiation of lake whitefish in northeastern Lake Michigan with implications for their management. *North Am. J. Fish. Manage.* **5**: 526-536.
- Schneeberger, P.J., Ebener, M.P., Toneys, M.L., and Peters, P.J. 2005a. Status of lake whitefish (*Coregonus clupeaformis*) in Lake Michigan. *In* Proceedings of a workshop on the dynamics of lake whitefish (*Coregonus clupeaformis*) and the amphipod *Diporeia* spp. in the Great Lakes. *Edited by* L.C. Mohr and T.F. Nalepa, Great Lakes Fish. Comm. Tech. Rep. No. 66. pp. 67-86.

- Schneeberger, P.J., Elliott, R.F., Jonas, J.L., and Hart, S. 2005b. Benthivores. *In* The state of Lake Michigan in 2000. *Edited by* M.E. Holeý and T.N. Trudeau. Great Lakes Fish. Comm. Spec. Pub. 05-01. pp. 59-73.
- Schneeberger, P.J., Toneys, M., Elliott, R., Jonas, J., Clapp, D., Hess, R., and Passino-Reader, D. 1998. Lakewide assessment plan for Lake Michigan fish communities [online]. Available from <http://www.glf.org/pubs/SpecialPubs/lwasses01.pdf> [accessed 2 Sept 2008].
- Scribner, K.T., Bott, K., Forsythe, P., and DeHaan, P. 2004. Lake Michigan Basin genetics sub-project: 2004 final report. Final report for the Great Lakes Fishery Trust. Proj. No. 2001.113. Mich. State Univ., Dept. Nat. Resour., East Lansing, MI.
- Seelbach, P.W. 1985. Smolt migration of wild and hatchery-raised coho and Chinook salmon in a tributary of northern Lake Michigan. Mich. Dept. Nat. Resour., Fish. Div., Fish. Res. Rep. No. 1935.
- Seelbach, P.W. 1987a. Smolting success of hatchery-raised steelhead planted in a Michigan tributary of Lake Michigan. *N. Am. J. Fish. Manage.* **7**: 223-231.
- Seelbach, P.W. 1987b. Effect of winter severity on steelhead smolt yield in Michigan: an example of the importance of environmental factors in determining smolt yield. *Am. Fish. Soc. Symp.* **1**: 441-450.
- Seelbach, P.W., Dexter, J.L., and Ledet, N.D. 1994. Performance of steelhead smolts stocked in southern Michigan warmwater rivers. Mich. Dept. Nat. Resour., Fish. Div., Fish. Res. Rep. No. 2003.
- Seelbach, P.W., and Whelan, G.E. 1988. Identification and contribution of wild and hatchery steelhead stocks in Lake Michigan tributaries. Mich. Dept. Nat. Resour., Fish. Div., Fish. Res. Rep. No. 1950.
- Seelbach, P.W., and M.J. Wiley. 2005. An initial, landscape-based information system for ecological assessment of Lake Michigan tributaries. *In* State of Lake Michigan: ecology, health and management. *Edited by* T. Edsall and M. Munawar. Ecovision World Mono. Ser., Aquatic Ecosystem Health Manage. Soc. pp. 559-581.
- Shroyer, S.M., and McComish, T.S. 1998. Forecasting abundance of quality-sized yellow perch in Indiana waters of Lake Michigan. *North Am. J. Fish. Manage.* **18**: 19-24.
- Shroyer, S.M., and McComish, T.S. 2000. Relationship between alewife abundance and yellow perch recruitment in southern Lake Michigan. *North Am. J. Fish. Manage.* **20**: 220-225.

- Simon, T.P., Dufour, R.L. Stewart, P.M., and Moffett, M.L. 2005. Distribution, classification and selection rationale of coastal wetlands in Lake Michigan. *In* State of Lake Michigan: Ecology, health and management. *Edited by* T. Edsall and M. Munawar. Ecovision World Mono. Ser., Aquatic Ecosystem Health Manage. Soc. pp. 479-502.
- Smith, S.H. 1970. Species interactions of the alewife in the Great Lakes. *Trans. Am. Fish. Soc.* **99**: 754-765.
- Sprules, W.G., Jin, E.H., Herman, A.W., and Stockwell, J.D. 1998. Calibration of an optical plankton counter for use in fresh water. *Limnol. Oceanogr.* **43**(4): 726-733.
- Steinhart, G.B., Stein, R.A., and Marschall, E.A. 2004. High growth rate of young-of-the-year smallmouth bass in Lake Erie: a result of the round goby invasion? *J. Great Lakes Res.* **30**: 381-391.
- Stephen, C., and Thorburn, M. 2002. Formulating a vision for fish health research in the Great Lakes [online]. Available from <http://www.glfrc.org/research/reports/stephen_fish_health.pdf> [accessed 9 September 2008].
- Stewart, D.J., and Binkowski, F.P. 1986. Dynamics of consumption and food conversion by Lake Michigan alewives: an energetics-modeling synthesis. *Trans. Am. Fish. Soc.* **115**: 643-661.
- Stewart, D.J., Kitchell, J.F., and Crowder, L.B. 1981. Forage fish and their salmonid predators in Lake Michigan. *Trans. Am. Fish. Soc.* **110**: 751-763.
- Swank, D. 2005. Life-history variation and management of wild Great Lakes steelhead populations. Ph.D. thesis. Univ. of Mich., Ann Arbor, MI.
- Szalai, E.B. 2003. Uncertainty in the population dynamics of alewife (*Alosa pseudoharengus*) and bloater (*Coregonus hoyi*) and its effects on salmonine stocking strategies in Lake Michigan. Ph.D. thesis. Mich. State Univ., East Lansing, MI.
- Taube, C.M. 1974. Transfer releases of coho salmon and trout into an upper part of Platte River, and observations on salmonid spawning. Mich. Dept. Nat. Resour., Fish. Div., Fish. Res. Rep. No. 1815.
- Thuemler, T.F. 1997. Lake sturgeon management in the Menominee River, a Wisconsin-Michigan boundary water. *Environ. Biol. Fishes* **48**: 311-317.
- Tillitt, D.E., Zajicek, J., Brown, S.B., Brown, L., Fitzsimons, J.D., Honeyfield, D.C., Holey, M., and Wright, G. 2005. Thiamine and thiaminase status in forage fish of salmonines from Lake Michigan. *J. Aquat. Anim. Health.* **17**: 13-25.

- Trudeau, T.N. 2005. The state of Lake Michigan in 2000 and its linkage to the joint strategic plan for management of the Great Lakes. *In* State of Lake Michigan: ecology, health and management. *Edited by* T. Edsall and M. Munawar. Ecovision World Mono. Ser., Aquatic Ecosystem Health Manage. Soc. pp. 521-536.
- Truemper, H.A., and Lauer, T.E. 2005. Gape limitation and piscine prey size-selection by yellow perch in the extreme southern area of Lake Michigan, with emphasis on two exotic prey items. *J. Fish Biol.* **66**: 135-149.
- United States vs. State of Michigan. 2000. Consent Decree, August 8, 2000. U.S. District Court, Western District of Michigan, Southern Division. Case No. 2:73 CV 26.
- U.S. EPA. 2004a. Updated information on Great Lakes areas of concern [online]. Available from <http://www.glin.net/aocstrategyteam/documents/AOCs2004.pdf> [accessed 9 September 2008].
- U.S. EPA. 2004b. Lake Michigan lakewide management plan 2004 status report [online]. Available from <http://www.epa.gov/glnpo/lakemich/2004update/> [accessed 9 September 2008].
- U.S. EPA. 2005. Ecological classification of rivers for environmental assessment [online]. Available from <http://sitemaker.umich.edu/riverclassproject> [accessed 9 September 2008].
- U.S. EPA. 2006. Great Lakes Legacy Act. [online]. Available from <http://www.epa.gov/glnpo/sediment/legacy/index.html> [accessed 9 September 2008].
- U.S. Fish and Wildlife Service/Fisheries and Oceans Canada. 2005. Protocol for application of lampricides to streams with populations of young-of-year lake sturgeon (*Acipenser fulvescens*). Tech. Oper. Proc. 011.4A. Marquette MI and Sault Ste. Marie, ON.
- Uzarski, D.G., Burton, T.M., Cooper, M.J., Ingram, J.W., and Timmermans, S. 2005. Fish habitat use within and across wetland classes in coastal wetlands of the five Great Lakes: development of a fish-based index of biotic integrity. *J. Great Lakes Res.* **31**(Suppl. 1): 171-187.
- Vanderploeg, H.A., Nalepa, T.F., Jude, D.J., Mills, E.F., Holeck, K.T., Liebig, J.T., Grigorovich, I.A., and Ojaveer, H. 2001. Dispersal and emerging ecological impacts of Ponto-Caspian species in the Laurentian Great Lakes. *Can. J. Fish. Aquat. Sci.* **59**:1209-1228.
- Walker, S.H., Prout, M.W., Taylor, W.W., and Winterstein, S.R. 1993. Population dynamics and management of lake whitefish stocks in Grand Traverse Bay, Lake Michigan. *North Am. J. Fish. Manage.* **13**: 73-85.

- Warner, D.M., Claramunt, R.M., Faul, C.S., and O'Brien, T.P. 2005. Status of pelagic prey fish in Lake Michigan, 2001-2004. *In* Minutes of the 2005 Annual Meeting of the Lake Michigan Committee. Great Lakes Fishery Commission, Ann Arbor, MI.
- Warren, G.J., and R.G. Kreis. 2005. Recent and long-term nutrient trends in Lake Michigan. *In* State of Lake Michigan: ecology, health and management. Edited by T. Edsall and M. Munawar. Ecovision World Mono. Ser., Aquatic Ecosystem Health Manage. Soc. pp. 141-155.
- Wei, A., Chow-Fraser, P., and Albert, D. 2004. Influence of shoreline features on fish distribution in the Laurentian Great Lakes. *Can. J. Fish. Aquat. Sci.* **61**(7): 1113-1123.
- Wells, L. 1980. Food of alewives *Alosa pseudoharengus*, yellow perch *Perca flavescens*, spottail shiners *Notropis hudsonius*, trout perch *Percopsis omiscomaycus*, and slimy *Cottus bairdi* and fourhorn sculpins *Myoxocephalus quadricornis* in southeastern Lake Michigan USA. *U.S. Fish Wildl. Serv. Tech. Papers* **98**: 1-12.
- Wells, L., and McLain, A.L. 1973. Lake Michigan—man's effects on native fish stocks and other biota. Great Lakes Fish. Comm. Tech. Rep. 20.
- Wilberg, M., Bence, J.R., Eggold, B.T., Makauskas, D., Clapp, D.F. 2005. Yellow perch dynamics in southwestern Lake Michigan during 1986-2002. *North Am. J. Fish. Manage.* **25**: 1130-1152.
- Wilcox, D.A. 2005. Lake Michigan wetlands: classification, concerns and management opportunities. *In* State of Lake Michigan: ecology, health and management. Edited by T. Edsall and M. Munawar. Ecovision World Mono. Ser., Aquatic Ecosystem Health Manage. Soc. pp. 421-437.
- Wiley, M.J. 2005. Muskegon River mega model [online]. Available from <http://148.61.56.211/megaindex.htm> [accessed 12 September 2008].
- Wisconsin Department of Natural Resources. 2000. Wisconsin's lake sturgeon management plan. Wisc. Dept. Nat. Resour. Bur. Fish. Manage. Habitat Protection.
- Witt, A.M., Dettmers, J.M., and Cáceres, C.E. 2005. *Cercopagis pengoi* in southwestern Lake Michigan in the four years following invasion. *J. Great Lakes Res.* **31**: 245-252.
- Woldt, A.P., and Rutherford, E.S. 2002. Production of juvenile steelhead in two central Lake Michigan tributaries. Mich. Dept. Nat. Resour., Fish. Div., Fish. Res. Rep. No. 2060.

- Woldt, A.P., Sitar, S.P., Bence, J.R., and Ebener, M.P (EDS.). 2004. Summary status of lake trout and lake whitefish populations in the 1836 treaty-ceded waters of lakes Superior, Huron and Michigan in 2002, with recommended yield and effort levels for 2003. Technical Fisheries Committee, 1836 treaty-ceded waters of Lakes Superior, Huron and Michigan.
- Woldt, A.P., Sitar, S.P., Bence, J.R., and Ebener, M.P (EDS.). 2005. Summary status of lake trout and lake whitefish populations in the 1836 treaty-ceded waters of lakes Superior, Huron and Michigan in 2003, with recommended yield and effort levels for 2004. [online]. Available from michigan.gov/documents/2004StatusLT&LakeWhitefishPop_126360_7.pdf [accessed 12 September 2008].
- Wright, G., Honeyfield, D.C., and Brown, S.B. 2005. Fish health. *In* The state of Lake Michigan in 2000. *Edited by* M.E. Holey and T.N. Trudeau. Great Lakes Fish. Comm. Spec. Pub. 05-01. pp. 81-86.
- Wurster, C.M., Patterson, W.P., Stewart, D.J., Bowlby, J.N., and Stewart, T.J. 2005. Thermal histories, stress, and metabolic rates of Chinook salmon (*Oncorhynchus tshawytscha*) in Lake Ontario: evidence from intra-otolith stable isotope analyses Can. J. Fish. Aquat. Sci. **62**(3): 700-713.
- Zafft, D.J. 1992. Migration of wild Chinook and coho salmon smolts from the Pere Marquette River, Michigan. M.Sc. thesis, Mich. State Univ., East Lansing, MI.
- Zorn, T.G., Seelbach, P.W., and Wiley, M.J. 2002. Distributions of stream fishes and their relationship to stream size and hydrology in Michigan's Lower Peninsula. Trans. Am. Fish. Soc. **131**: 70-85.

- 88-2 A proposal for a bioassay procedure to assess impact of habitat conditions on lake trout reproduction in the Great Lakes (report of the ad hoc Committee to Assess the Feasibility of Conducting Lake Trout Habitat Degradation Research in the Great Lakes). 1988. Edited by R. L. Eshenroder. 13 p.
- 88-3 Age structured stock assessment of Lake Erie walleye (report of the July 22-24, 1986 Workshop). 1988. R. B. Deriso, S. J. Nepszy, and M. R. Rawson. 13 p.
- 88-4 The international Great Lakes sport fishery of 1980. 1988. D. R. Talhelm. 70 p.
- 89-1 A decision support system for the integrated management of sea lamprey. 1989. J. F. Koonce and A. B. Locci-Hernandez. 74 p.
- 90-1 Fish community objectives for Lake Superior. 1990. Edited by T. R. Busiahn. 24 p.
- 90-2 International position statement and evaluation guidelines for artificial reefs in the Great Lakes. 1990. Edited by J. E. Gannon. 24 p.
- 90-3 Lake Superior: the state of the lake in 1989. 1990. Edited by M. J. Hansen. 56 p.
- 90-4 An ecosystem approach to the integrity of the Great Lakes in turbulent times (proceedings of a 1988 workshop supported by the Great Lakes Fishery Commission and the Science Advisory Board of the International Joint Commission). 1990. Edited by C. J. Edwards and H. A. Regier. 302 p.
- 91-1 Status of walleye in the Great Lakes: case studies prepared for the 1989 workshop. 1991. Edited by P. J. Colby, C. A. Lewis, and R. L. Eshenroder. 222 p.
- 91-2 Lake Michigan: an ecosystem approach for remediation of critical pollutants and management of fish communities (report of a round table sponsored in 1990 by the Great Lakes Fishery Commission, the Science Advisory Board of the International Joint Commission, and the Lake Michigan Federation). 1991. Edited by R. L. Eshenroder, J. H. Hartig, and J. E. Gannon. 58 p.
- 91-3 The state of the Lake Ontario fish community in 1989. 1991. S. J. Kerr and G. C. LeTendre. 38 p.
- 93-1 Great Lakes fish disease control policy and model program. 1993. Edited by J. G. Hnath. 38 p. Protocol to minimize the risk of introducing emergency disease agents with importation of salmonid fishes from enzootic areas. 1993. Edited by R.W. Horner and R. L. Eshenroder. 15 p.
- 94-1 The state of Lake Superior in 1992. 1994. Edited by M. J. Hansen. 110 p.
- 94-2 An introduction to economic valuation principles for fisheries management. L. G. Anderson. 98 p.
- 95-1 Fish-community objectives for Lake Huron. 1995. R. L. DesJardine, T. K. Gorenflo, R. N. Payne, and J. D. Schrouder. 38 p.
- 95-2 The state of Lake Huron in 1992. Edited by M. P. Ebener. 140 p.
- 95-3 Fish-community objectives for Lake Michigan. R.L. Eshenroder, M.E. Holey, T.K. Gorenflo, and R.D. Clark, Jr. 56 p.
- 99-1 Fish-community objectives for Lake Ontario. T.J. Stewart, R.E. Lange, S.D. Orsatti, C.P. Schneider, A. Mathers, M.E. Daniels. 56 p.
- 03-01 Fish-community objectives for Lake Superior. W.H. Horns, C.R. Bronte, T.R. Busiahn, M.P. Ebener, R.L. Eshenroder, T. Gorenflo, N. Kmiecik, W. Mattes, J.W. Peck, M. Petzold, D.R. Schreiner. 86 p.
- 03-2 Fish-community goals and objectives for Lake Erie. P.A. Ryan, R. Knight, R. MacGregor, G. Towns, R. Hoopes, W. Culligan. 56 p.
- 05-01 The state of Lake Michigan in 2000. Edited by M.E. Holey and T.N. Trudeau. 114 p.
- 05-02 The state of Lake Huron in 1999. Edited by M.P. Ebener. 140 p.
- 07-01 The state of Lake Ontario in 2003. Edited by B.J. Morrison and S. R. LaPan. 108 p.
- 07-02 The state of Lake Superior in 2000. Edited by M.P. Ebener. 126 p.
- 08-01 The state of Lake Huron in 2004. Edited by J.R. Bence and L.C. Mohr. 88 p.

Special Publications

- 79-1 Illustrated field guide for the classification of sea lamprey attack marks on Great Lakes lake trout. 1979. E. L. King and T. A. Edsall. 41 p.
- 82-1 Recommendations for freshwater fisheries research and management from the Stock Concept Symposium (STOCS). 1982. A. H. Berst and G. R. Spangler. 24 p.
- 82-2 A review of the adaptive management workshop addressing salmonid/lamprey management in the Great Lakes. 1982. Edited by J. F. Koonce, L. Greig, B. Henderson, D. Jester, K. Minns, and G. Spangler. 58 p.
- 82-3 Identification of larval fishes of the Great Lakes basin with emphasis on the Lake Michigan drainage. 1982. Edited by N. A. Auer. 744 p. (Cost: \$10.50 U.S., \$12.50 CAN)
- 83-1 Quota management of Lake Erie fisheries. 1983. Edited by J. F. Koonce, D. Jester, B. Henderson, R. Hatch, and M. Jones. 40 p.
- 83-2 A guide to integrated fish health management in the Great Lakes basin. 1983. Edited by F. P. Meyer, J. W. Warren, and T. G. Carey. 262 p.
- 84-1 Recommendations for standardizing the reporting of sea lamprey marking data. 1984. R. L. Eshenroder and J. F. Koonce. 22 p.
- 84-2 Working papers developed at the August 1983 conference on lake trout research. 1984. Edited by R. L. Eshenroder, T. P. Poe, and C. H. Olver.
- 84-3 Analysis of the response to the use of "Adaptive Environmental Assessment Methodology" by the Great Lakes Fishery Commission. 1984. C. K. Minns, J. M. Cooley, and J. E. Forney. 22 p.
- 85-1 Lake Erie fish community workshop (report of the April 4-5, 1979 meeting). 1985. Edited by J. R. Paine and R. B. Kenyon. 58 p.
- 85-2 A workshop concerning the application of integrated pest management (IPM) to sea lamprey control in the Great Lakes. 1985. Edited by G. R. Spangler and L. D. Jacobson. 98 p.
- 85-3 Presented papers from the Council of Lake Committees plenary session on Great Lakes predator-prey issues, March 20, 1985. 1985. Edited by R. L. Eshenroder. 134 p.
- 85-4 Great Lakes fish disease control policy and model program. 1985. Edited by J. G. Hnath. 24 p.
- 85-5 Great Lakes Law Enforcement/Fisheries Management Workshop (report of the 21, 22 September 1983 meeting). 1985. Edited by W. L. Hartman and M. A. Ross. 26 p.
- 85-6 TFM vs. the sea lamprey: a generation later. 1985. 18 p.
- 86-1 The lake trout rehabilitation model: program documentation. 1986. C. J. Walters, L. D. Jacobson, and G. R. Spangler. 34 p.
- 87-1 Guidelines for fish habitat management and planning in the Great Lakes (report of the Habitat Planning and Management Task Force and Habitat Advisory Board of the Great Lakes Fishery Commission). 1987. 16 p.
- 87-2 Workshop to evaluate sea lamprey populations "WESLP" (background papers and proceedings of the August 1985 workshop). 1987. Edited by B. G. H. Johnson.
- 87-3 Temperature relationships of Great Lakes fishes: a data compilation. 1987. D. A. Wismer and A. E. Christie. 196 p.
- 88-1 Committee of the Whole workshop on implementation of the Joint Strategic Plan for Management of Great Lakes Fisheries (reports and recommendations from the 18-20 February 1986 and 5-6 May 1986 meetings). 1988. Edited by M. R. Dochoda. 170 p.

